

**BIRDS of the upper Mazar watershed**

Compiled by the Fundación Cordillera Tropical, January 2011

Order/Family	Species	Common name (English)	Nombre común (español)	Habitat	Population	Status	Comments
<b>ANSERIFORMES</b>							
Anatidae	<i>Anas andium</i>	Andean teal	Cerceta andina	L	uncommon		
	<i>Oxyura ferruginea</i>	Andean duck	Pato rojizo andino	L	uncommon		
	<i>Merganetta armata</i>	Torrent duck	Pato torrentero	R	uncommon		
<b>APODIFORMES</b>							
Apodidae	<i>Streptoprocne zonaris</i>	White-collared swift	Vencejo cuelliblanco	P	common		
	<i>Cypseloides rutilus</i>	Chestnut-collared swift	Vencejo cuellicastañó	F, CL	uncommon		
Trochilidae	<i>Aglaeactis cupripennis</i>	Shining sunbeam	Rayito brillante	F	common		Seen one at the treeline in these flowers.
	<i>Aglaiocercus kingi</i>	Long-tailed sylph	Silfo colilargo	F, CL	uncommon		
	<i>Chalcostigma herrani</i>	Rainbow-bearded thornbill	Picoespina arcoiris	F	uncommon		Seen along a primary forest path with <i>Chusquea</i> bamboo.
	<i>Coeligena iris</i>	Rainbow starfrontlet	Frentiestrella arcoiris	F	common		Rather common species, rarely seen well. Feeds on bromeliads.
	<i>Coeligena lutetiae</i>	Buff-winged starfrontlet	Frentiestrella alianteada				Not previously recorded in Mazar. One recorded from treeline shrub.
	<i>Colibri coruscans</i>	Sparkling violetear	Orejivioleta ventriazul	F	common		
	<i>Ensifera ensifera</i>	Sword-billed hummingbird	Colibrí pico espada	F	uncommon		One seen along a stream at the bottom of the dirt road, holding its long bill upward. It dashed away quickly.
	<i>Eriocnemis luciani</i>	Sapphire-vented puffleg	Zamarrito colilargo	F	common		
	<i>Eriocnemis vestitus</i>	Glowing puffleg	Zamarrito luciente	F	uncommon		Seen once on the treeline, in low bushes.
	<i>Helianthus exortis</i>	Tourmaline sunangel	Solángel turmalina	F	uncommon		Not previously recorded at Mazar. A female with conspicuous white throat was seen in primary forest. Species distinction between <i>H. exortis</i> and <i>H. micraster</i> is not universally accepted; since observations of both were made within 10 minutes in the same location, it's possible that they belonged to the same species.
	<i>H. micraster</i>	Flame-throated sunangel	Solángel gorjidorado				Seen once in primary forest, close to the same spot as <i>H. exortis</i> (see discussion above). Orange on chest visible from about 50m.
	<i>H. viola</i>	Purple-throated sunangel	Solángel gorjipúrpura	F	common		Uncommon near bigger streams and in eucalyptus trees in pastures.
	<i>Lafresnaya lafresnaya</i>	Mountain velvetbreast	Colibrí terciopelo	F	uncommon		A common and aggressive hummer of the flowers, often hiding under a little roof. Otherwise only seen along forest edge and lower scrubs. Fast swaying flight with white tail distinctive.
	<i>Lesbia nuna</i>	Green-tailed trainbearer	Colacinta coliverde	P	uncommon		A female constantly present in a flowering bush.
	<i>L. victoriae</i>	Black-tailed trainbearer	Colacinta colinegra		uncommon		Seen once at the bottom of a dirt road along the river.
	<i>Metalura baroni</i>	Violet-throated metaltail	Metalura gorjivioleta	P	uncommon, endemic		
	<i>M. tyrianthina</i>	Tyrian metaltail	Metalura tiria	F	common		In pastures and secondary forest, often in Eucalyptus trees.
	<i>M. williami</i>	Viridian metaltail	Metalura verde	P	common		
	<i>Patagona gigas</i>	Giant hummingbird	Colibrí gigante	CL	common		
	<i>Phaethornis syrmatorphorus</i>	Tawny-bellied hermit	Ermitaño ventrileonado	F			
	<i>Pterophanes cyanopterus</i>	Great sapphirewing	Alizafiro grande		uncommon		Not previously recorded in Mazar. Observed twice right at the treeline, and at one spot 300 meters below, feeding on pink flowers.
<b>CAPRIMULGIFORMES</b>							
Nyctibiidae	<i>Nyctibius maculosus</i>	Andean potoo	Nictibio andino		rare		Recording made of this rare and local species. Probably responsible for the dead large beetles found in and below a trunk in pasture?
Caprimulgidae	<i>Caprimulgus longirostris</i>	Band-winged nightjar	Chotacabras alifajeado	F	uncommon		Often sitting on the dirt road at night. We flushed one from a little fence post right forest edge at dawn.
<b>CHARADRIFORMES</b>							
Charadriidae	<i>Vanellus resplendens</i>	Andean lapwing	Avefría andina	L	uncommon		
Laridae	<i>Larus serranus</i>	Andean gull	Gaviota andina	CL	uncommon		Uncommon, seen twice above the valley.
Scolopacidae	<i>Actitis macularia</i>	Spotted sandpiper	Andarríos coleador	L	uncommon		
	<i>Tringa melanoleuca</i>	Greater yellowlegs	Patiamarillo mayor	L	uncommon		
	<i>Gallinago jamesoni</i>	Andean snipe	Becasina andina	L	uncommon		
	<i>G. imperialis</i>	Imperial snipe	Becasina imperial	F	rare		Unconfirmed

Order/Family	Species	Common name (English)	Nombre común (español)	Habitat	Population	Status	Comments
<b>CICONIIFORMES</b> Cathartidae	<i>Coragyps atratus</i> <i>Cathartes aura</i>	Black vulture Turkey vulture	Gallinazo negro Gallinazo cabecirrojo	CL F, CL, P	uncommon rare		One seen over the valley. More common near Zhoray. Seen twice over the valley on sunny days.
<b>COLUMBIFORMES</b> Columbidae	<i>Claravis mondetoura</i> <i>Columba fasciata</i> <i>Leptotila verreauxi</i>	Maroon-chested ground-dove Band-tailed pigeon White-tipped dove	Tortolita pechimarrón Paloma collaraja Paloma apical	 F F	 uncommon uncommon		Not previously recorded in Mazar. One hypothetically recorded. Rather shy, seen only occasionally in small groups over the canopy. A rather common dove on the pastures below the forest.
<b>FALCONIFORMES</b> Accipitridae	<i>Accipiter ventralis</i> <i>Buteo leucorrhous</i> <i>Buteo polyosoma</i> <i>Circus cinereus</i> <i>Elanoides forficatus</i> <i>Geranoaetus melanoleucus</i> <i>Oroaetus isidori</i>	Plain-breasted hawk White-rumped hawk Variable hawk Cinereous harrier Swallow-tailed kite Black-chested buzzard-eagle Black-and-chestnut eagle	Azor pechillano Gavilán lomiblanco Gavilán variable Aguilucho cinéreo Elanio tijereta Aguila pechinegra Aguila andina	F, P F, P F, CL, P P P F, CL, P F	uncommon rare uncommon rare uncommon	Ap. 2      near threatened	One seen next to a dirt road.  One seen over the valley on a sunny day.   Twice, a couple soared for long periods of time over the valley. Not previously recorded at Mazar. A juvenile sighted soaring at just a few meters distance, being chased by a Kestrel. Not previously recorded in Mazar.
Falconidae	<i>Falco femoralis</i> <i>F. peregrinus</i> <i>F. sparverius</i> <i>Phalcoboenus carunculatus</i>	Aplomado falcon Peregrine falcon American kestrel Carunculated caracara	Halcón aplomado Halcón peregrino Cernícalo americano Caracara curiquingue	 P F, CL, P P	rare rare common uncommon		Territorial, displaying great stamina in getting rid of other raptors.
<b>GALLIFORMES</b> Cracidae	<i>Penelope montagnii</i>	Andean guan	Pava andina	F	common		Rather common, and noisy in primary forest up to the treeline.
<b>PASSERIFORMES</b> Cardinalidae	<i>Pheucticus chrysogaster</i> <i>P. aureoventris</i> <i>Saltator cinctus</i>	Southern yellow-grosbeak Black-backed grosbeak Masked saltator	Picogrueso amarillo sureño Picogrueso dorsinegro Saltador enmascarado	CL F F	common uncommon rare	near threatened	Seen twice in secondary growth bushes and pastures.
Cinclidae	<i>Cinclus leucocephalus</i>	White-capped dipper	Cinco gorriblanco	R	uncommon		One seen along large river.
Corvidae	<i>Cyanolyca turcosa</i>	Turquoise jay	Urraquita turquesa	F, CL	common		Often roaming around in large groups with Northern Mountain Caciques. Makes many different sounds. Unconfirmed; calls recorded from primary forest near the treeline.
Cotingidae	<i>Ampelion rubrocristatus</i> <i>Pipreola arcuata</i>	Red-crested cotinga Barred fruiteater	Cotinga crestirroja Frutero barreteado	F F			Observed twice, one female and a pair; heard more often.
Dendrocolaptidae	<i>Campylorhamphus pucherani</i> <i>Dendrocincla tyrannina</i> <i>Lepidocolaptes affinis</i> <i>L. lacrymiger</i> <i>Xiphocolaptes promeropirhynchus</i>	Greater scythebill Tyrannine woodcreeper Spot-crowned woodcreeper Montane woodcreeper Strong-billed woodcreeper	Picoguadaña grande Trepatroncos tiranino  Trepatroncos montano Trepatroncos piquifuerte	F F F F	rare uncommon	near threatened	Formerly considered conspecific w/ <i>L. lacrymiger</i> . Uncommon in species flocks, where found singly or in pairs. Not previously recorded in Mazar. With an altitudinal range of 1,100 to 3,000m (Ridgely and Greenfield 2001), it is at its height limit here. Found practically every day in species flocks or alone in small groups of up to five individuals.
Emberizidae	<i>Atlapetes latinuchus</i> <i>A. pallidinucha</i> <i>A. schistaceus</i> <i>Buarremon torquatus</i> <i>Catamenia inornata</i> <i>Phrygilus unicolor</i> <i>Zonotrichia capensis</i> <i>Grallaria squamigera</i>	Rufous-naped brush-finch Pale-naped brush-finch Slatey brush-finch Stripe-headed brush finch Plain-colored seedeater Plumbeous sierra-finch Rufous-coloured sparrow Undulated antpitta	Matorralero nuquirrufo Matorralero nuquipálido Matorralero pizarroso Matorralero cabecilistado Semillero sencillo Frigilo plumizo Chingolo Gralaria ondulada	F F F F F P F, P, CL F	uncommon  uncommon uncommon common common uncommon common		Not previously recorded in Mazar. Seen once at treeline in a group with Rufous-naped, Bar-winged cinclodes and White-throated tyrannulet. Seen twice at the treeline with Rufous-naped. Fairly common in primary forest, sometimes with species flocks. Fairly common along secondary growth edges.
Formicariidae	<i>G. ruficapilla</i> <i>G. rufula</i>	Chestnut-crowned antpitta Rufous antpitta	Gralaria cornicastaña Gralaria rufa	 F	common uncommon		Common in cleared areas; none seen in paramo. Relatively easy to observe along primary forest paths at dawn and dusk. Also found right outside the primary forest. Not previously recorded in Mazar. Fairly common at forest edge, often heard, not seen. Heard throughout the day, from dawn till after dusk. Also seen once at dawn along a primary forest path, about 2m up on a branch.

Order/Family	Species	Common name (English)	Nombre común (español)	Habitat	Population	Status	Comments
<b>PASSERIFORMES</b>							
Formicariidae	<i>G. quitensis</i>	Tawny antpitta	Gralaria leonada	F, P	common		Fairly common at treeline, often heard, not seen.
	<i>Grallaricula lineifrons</i>	Crescent-faced antpitta	Gralarita carilunada	F	extremely rare	near threatened	
Furnariidae	<i>Asthenes flammulata</i>	Many-striped canastero	Canastero multilistado	P	uncommon		
	<i>Cinclodes excelsior</i>	Stout-billed cinclodes	Cinclodes piquigruoso		uncommon		
	<i>Cinclodes fuscus</i>	Bar-winged cinclodes	Cinclodes alifranjeado	P			Seen once on a misty day at the treeline, in a group with White-throated Tyrannulet, Pale-naped and Rufous-naped Brush-Finches.
	<i>Hellmayrea gularis</i>	White-browed Spinetail	Colaespina cejiblanca				
	<i>Margarornis squamiger</i>	Pearled treerunner	Subepalo perlado	F			Common in species flocks, often in small groups of three or four, sticking together. Usually seen in larger branches in the canopy.
	<i>Pseudocolaptes boissonneautii</i>	Streaked tuftedcheek	Barbablanca rayada	F			Uncommon visitor to species flocks, seen twice.
	<i>Schizoeaca griseomurina</i>	Mouse-coloured thistletail	Colicardo murino		uncommon		
	<i>Synallaxis azarae</i>	Azara's spinetail	Colaespina de Azara	F, CL			Common in secondary growth and pastures.
	<i>Thripadectes flammulatus</i>	Flammulated treehunter	Trepamusgos flamulado				Not previously recorded at Mazar. One hypothetical bird recorded.
Hirundinidae	<i>Notiochelidon murina</i>	Brown-bellied swallow	Golondrina ventricafé	F, P, CL	common		Usually seen in the morning and before dusk above pastures and the forest canopy. Nests in houses.
	<i>N. cyanoleuca</i>	Blue-and-white swallow	Golondrina azuliblanca	F, CL	rare		Not uncommonly with Brown-bellieds, also nesting in house.
	<i>N. flavipes</i>	Pale-footed swallow	Golondrina nubiselvática	F, CL	rare		
Icteridae	<i>Amblycercus holosericeus</i>	Yellow-billed cacique	Cacique piquiamarillo	F	unknown		
	<i>Cacicus leucoramphus</i>	Northern mountain-cacique	Cacique montañas norteño	F	common		Often in association with Turquoise Jays, and usually noisy.
	<i>Carduelis magellanica</i>	Hooded siskin	Jilguero encapuchado	CL	common		Two small groups seen high up in Eucalyptus trees.
	<i>Sturnella bellicosa</i>	Peruvian meadowlark	Pastorero peruano	CL	common		
	<i>S. militaris</i>	Red-breasted blackbird	Pastorero pechirrojo	CL	common		
Parulidae	<i>Anthus bogotensis</i>	Paramo pipit	Bisbita del páramo	P	uncommon		
	<i>Basileuterus coronatus</i>	Russet-crowned warbler	Reinita coronirrojiza	F	common		Common in primary forest, heard often during dawn chorus. Sometimes accompanies species flock.
	<i>B. luteoviridis</i>	Citrine warbler	Reinita citrina	F	uncommon		Uncommon, found three times, along forest edges and in primary forest, in presence of Black-crested Warblers.
	<i>B. nigrocristatus</i>	Black-crested warbler	Reinita crestinegra	F	common		Commonly heard during dawn chorus in primary forest, with a variety of songs. Also seen often with species flocks.
	<i>Dendroica fusca</i>	Blackburnian warbler	Reinita pechinaranja	F	common		
	<i>Myioborus melanocephalus</i>	Spectacled whitestart	Candelita de anteojos	F	common		The most common warbler. Found every day in all habitats.
	<i>M. miniatus</i>	Slate-throated whitestart	Candelita goliplomiza	CL			
Rhinocryptidae	<i>Acropternis orthonyx</i>	Ocellated tapaculo	Tapaculo ocelado	F	uncommon		
	<i>Myornis senilis</i>	Ash-colored tapaculo	Tapaculo cenizo	F	unknown		
	<i>Scytalopus canus</i>	Paramo tapaculo	Tapaculo paramero	P	unknown		Not previously recorded at Mazar. Heard practically every day.
	<i>Scytalopus unicolor</i>	Unicolored tapaculo	Tapaculo unicolor	F	uncommon		Uncommonly heard in primary forest.
Thraupidae	<i>Anisognathus igniventris</i>	Scarlet-bellied mountain-tanager	Tangara montana ventriescarlata	F	common		Surprisingly uncommon. Seen once at treeline with Black-chested Mountain-Tanagers, and once in a species flock with Buff-breasted and Lacrimose Mountain-Tanagers. Not found in pasture or secondary forest.
	<i>A. lacrymosus</i>	Lacrimose mountain-tanager	Tangara montana lagrimosa	F	common		Seen in species flocks. Usually silent and not conspicuous.
	<i>Buthraupis montana</i>	Hooded mountain-tanager	Tangara montana encapuchada	F, CL	uncommon		Fairly common in canopy of primary forest. Song sometimes the first to start the dawn chorus.
	<i>B. eximia</i>	Black-chested mountain-tanager	Tangara montana pechinegra	F	rare		Three birds at the treeline on a misty day with Scarlet-bellieds.
	<i>B. wetmorei</i>	Masked mountain-tanager	Tangara montana enmascarada	F	uncommon		
	<i>Catamblyrhynchus diadema</i>	Plushcap	Gorradidema	F	uncommon		Fairly common in and near <i>Chusquesa</i> bamboo thickets.
	<i>Cnemoscopus rubrirostris</i>	Grey-hooded bush-tanager	Tangara montés capuchigris	F	uncommon		
	<i>Conirostrum cinereum</i>	Cinerous conebill	Picocono cinéreo	F, CL	common		Found twice in secondary forest edge.
	<i>C. sitticolor</i>	Blue-backed conebill	Picocono dorsiazul	F	common		Found twice in species flocks, alone on both occasions.
	<i>Diglossa humeralis</i>	Black flowerpiercer	Pinchaflor negro	F	common		Seen once along river at the bottom of a dirt road, among low bushes on a steep cliff with lots of bromeliads.
	<i>D. lafresnayii</i>	Glossy flowerpiercer	Pinchaflor satinado	F	common		

Order/Family	Species	Common name (English)	Nombre común (español)	Habitat	Population	Status	Comments	
<b>PASSERIFORMES</b>								
Thraupidae	<i>Diglossopsis cyanea</i>	Masked flowerpiercer	Pinchaflor enmascarado	F	uncommon		A fairly common tanager from primary and secondary forests, usually noted by its song in these dense habitats.	
	<i>Diglossopsis caerulescens</i>	Bluish flowerpiercer	Pinchaflor azulado	F				
	<i>Dubusia taeniata</i>	Buff-breasted mountain-tanager	Tangara montana pechianteadá	F	rare		Regular sightings of up to three birds at a time in species flocks. Usually silent or just giving contact calls.	
	<i>Hemispingus atropileus</i>	Black-capped hemispingus	Hemispingo coroninegro	F	uncommon		Seen twice in species flocks.	
	<i>H. superciliaris</i>	Superciliaried hemispingus	Hemispingo superciliado	F	uncommon		The most common Hemispingus, seen often in association with Spectacled Whitestarts and Black-crested Warblers.	
	<i>H. verticalis</i>	Black-headed hemispingus	Hemispingo cabecinegro	F	uncommon		Seen twice, in primary forest, very briefly on both occasions.	
	<i>Iridosornis rufivertex</i>	Golden-crowned tanager	Tangara coronidorada	F	uncommon		A trio observed at the treeline on a misty and rainy day.	
	<i>Tangara vassorii</i>	Blue-and-black tanager	Tangara azulinegra	F	uncommon		Seen only once near treeline in primary forest.	
	<i>Urothraupis stolzmanni</i>	Black-backed bush-tanager	Quinero dorsinegro	F	unknown			
	Troglodytidae	<i>Cinnycerthia unirufa</i>	Rufous wren	Soterrey rufo	F	uncommon		Small groups in stands of <i>Chusquea</i> bamboo, often near streams.
		<i>Cistothorus platensis</i>	Grass wren	Soterrey sabanero	P	uncommon		
		<i>Thryothorus euophrys</i>	Plain-tailed wren	Soterrey colillano	F	uncommon		Heard throughout the day; observed once in a large tract of bamboo.
<i>Troglodytes solstitialis</i>		Mountain wren	Soterrey montañés	F	common		Uncommon, found singly or in pairs, sometimes with species flocks.	
<i>Henicorhina leucophrys</i>		Gray-breasted wood-wren	Soterrey-montés pechigris		common			
<i>Turdus fuscater</i>		Great thrush	Mirlo grande	F, P, CL	common		Seen in pastures and secondary growth, but also in primary forest, even following species flocks at one occasion.	
Turdidae	<i>T. serranus</i>	Glossy-black thrush	Mirlo negribriloso	CL	uncommon			
	<i>Agriornis montana</i>	Black-billed shrike-tyrant	Arriero piquinegro	P	uncommon			
	<i>Anairetes parulus</i>	Tufted tit-tyrant	Cachudito torito	F, P	rare		Seen once in secondary growth along with Cinereous Conebill and Bar-bellied Woodpecker, and again at the treeline in some bushes.	
	<i>Chenarchus erythropygius</i>	Red-rumped bush-tyrant	Alinaranja lomirrojiza	P	uncommon			
	<i>Contopus fumigatus</i>	Smoke-colored pewee	Pibí ahumado	F	common			
	<i>Elaenia pallatangae</i>	Sierran elaenia	Elenia serrana	P			One in a species flock, in primary forest.	
	<i>Hemitriccus rufigularis</i>	Buff-throated tody-tyrant	Tirano todi golianteado				Not previously recorded in Mazar. Seen twice in species flocks, from close range; in rather open shrubs and trees in primary forest.	
	<i>Mecocerculus leucophrys</i>	White-throated tyrannulet	Tiranillo barbiblanco	F	common		One at the treeline, with Bar-winged Cinclodes and Pale-naped and Rufous-naped Brush-Finches.	
	<i>M. stictopterus</i>	White-banded tyrannulet	Tiranillo albibandeado	F	uncommon		Common in species flocks, and often heard singly in the forest.	
	<i>Myiotheretes fumigatus</i>	Smoky bush-tyrant	Alinaranja ahumada	F	uncommon		Seen once along forest edge at lower limit of the forest.	
	<i>M. striaticollis</i>	Streak-throated bush-tyrant	Alinaranja golilistada	F, CL	uncommon			
	<i>Ochthoeca cinnamomeiventris</i>	Slaty-backed chat-tyrant	Pitajo dorsipizarro	F, R	uncommon			
	<i>O. diadema</i>	Yellow-bellied chat-tyrant	Pitajo ventriamarillo	F	uncommon			
	<i>O. frontalis</i>	Crowned chat-tyrant	Pitajo coronado	P	rare			
	<i>O. fumicolor</i>	Brown-backed chat-tyrant	Pitajo dorsipardo	F, P	rare			
	<i>O. rufipectoralis</i>	Rufous-breasted chat-tyrant	Pitajo pechirrufo	F	uncommon		Found in secondary growth, and sometimes forest. Recorded a short chuk-chuk call (not on CDs); also a more explosive chatter.	
	<i>Pachyrampus versicolor</i>	Barred becard	Cabezón barreteado	F			Not seen, but recorded twice in species flocks in primary forest.	
	<i>Pseudotriccus ruficeps</i>	Rufous-headed pygmy-tyrant	Tirano enano cabecirrufo				Not previously recorded in Mazar. Observed on three occasions, singly or in pairs, scurrying low through the undergrowth.	
<i>Pyrrhomyias cinnamomea</i>	Cinnamon flycatcher	Mosquerito canelo	F	common		Often in association with species flocks but also found singly.		
<i>Sayornis nigricans</i>	Black phoebe	Febe guardarríos	R	uncommon				
<i>Serpophaga cinerea</i>	Torrent tyrannulet	Tiranolete guardarríos	R	rare				
<i>Uromyias agilis</i>	Agile tit-tyrant	Cachudito agil	F	uncommon				
<b>PICIFORMES</b>								
Picidae	<i>Campephilus pollens</i>	Powerful woodpecker	Carpintero poderoso				Not previously recorded in Mazar. Recorded during dawn chorus, at forest edge next to tract of <i>Chusquea</i> bamboo at 3,000m (usually encountered at 1,700-2,600m--Ridgely and Greenfield 2001).	
	<i>Piculus rivolii</i>	Crimson-mantled woodpecker	Carpintero dorsicarmesí	F	uncommon		Observed twice: one along forest edge with adjacent secondary growth, and one in primary forest.	

Order/Family	Species	Common name (English)	Nombre común (español)	Habitat	Population	Status	Comments
<b>PICIFORMES</b>							
Picidae	<i>Veniliornis nigriceps</i>	Bar-bellied woodpecker	Carpintero ventribarrado	F	uncommon		Seen in secondary growth bushes.
Ramphastidae	<i>Andigena hypoglauca</i>	Grey-breasted mountain-toucan	Tucán andino pechigris	F	common	near threatened	Heard a few times, but seen only twice. Once at dusk, flying over the canopy, and once at dawn, eating fruits in a tree.
<b>PSITTACIFORMES</b>							
Psittacidae	<i>Amazona mercenaria</i>	Scaly-naped parrot	Amazona nuquiescamosa	F	uncommon	endangered	
	<i>Hapalopsittaca pyrrhops</i>	Red-faced parrot	Loro carrirojo	F	uncommon	vulnerable	
	<i>Leptosittaca branickii</i>	Golden-plumed parakeet	Perico cachetidorado	F	rare		
	<i>Pionus seniloides</i>	White-capped parrot	Loro gorriblanco	F	uncommon		Only parrot species seen, observed daily in groups of up to 40.
	<i>Touit stictoptera</i>	Spot-winged parrotlet	Periquito alipunteado	F	uncommon		
<b>STRIGIFORMES</b>							
Tytonidae	<i>Tyto alba</i>	Barn owl	Lechuza campanaria	CL	uncommon		
Strigidae	<i>Asio stygius</i>	Stygian owl	Búho estigio	F	rare		
	<i>Bubo virginianus</i>	Great horned owl	Búho coronado americano	F	uncommon		
	<i>Glaucidium jardinii</i>	Andean pygmy owl	Mochuelo andino	F	uncommon		
	<i>Otus albobularis</i>	White-throated screech-owl	Autillo goliblanco	F	uncommon		Recorded one from forest edge on our last day.
	<i>Strix albitarsus</i>	Rufous-banded owl	Búho rufibandeado				Not previously recorded in Mazar. Heard often, two vocalisations.
<b>TINAMIFORMES</b>							
Tinamidae	<i>Nothocercus julius</i>	Tawny-breasted tinamou	Tinamú pechileonado		rare		Seen on two occasions near streams, walking over the path.
<b>TROGONIFORMES</b>							
Trogonidae	<i>Pharomachrus antisianus</i>	Crested quetzal	Quetzal crestado				Not previously recorded in Mazar. Observed in primary forest, flashing its white tail feathers.
	<i>Pharomachrus auriceps</i>	Golden-headed quetzal	Quetzal cabecidorado	F	uncommon		
	<i>Trogon personatus</i>	Masked trogon	Trogon personatus	F	uncommon		Recorded sound a few times at dawn.

<b>Habitat:</b>				
F- Forest	R- Riverine	L- Lakes	P- Paramo	CL- Cultivated land

**From combined inventories of B. Planqué, R. Ridgely, M. Hancock, V. Egan, F. Toral, B. Trujillo, and M. Pacheco, and Round River Conservation Studies (RRCS), in upper Mazar and neighboring watersheds**

Hancock, Mark, Veronica Egan, Fabian Toral and Beatrix Trujillo, 1989, "Birds of montane forest and paramo, Mazar and Juval River Catchmentes, Bosque Protector 15, Cañar, Ecuador," unpublished inventory.

Pacheco, Milton Xavier, y Edison Gerardo Correa. 2002. Lista de las principales especies inventariadas por los pobladores de la comunidad Queseras y Dudas, durante enero del 2001 a diciembre del 2002, en bosques nativos, páramos, y tierras de cultivo.

Proyecto Conservación y manejo del páramo en la comunidad Queseras, Fundación Ecohomode.

Planqué, Bob, 2002, "Bird Report from the Mazar Wildlife Reserve, Ecuador", <http://www.few.vu.nl/~rplanque/Mazar/mazar-bird-report.html>; see also Xeno-Canto song collection at <http://www.xeno-canto.org/browse.php?query=rec%3Aplanqu+loc%3Alibertad>

Ridgely, Robert, Percy Stafford, Rick Marsi and Francisco Sornoza, 1991, "Bird list of the Hacienda La Libertad," unpublished inventory.

Ridgely, Robert and Paul Greenfield, 2001, Birds of Ecuador: Field Guide (Cornell University Press, Ithaca).